

hep-ph/9707468

(SEMI)CLASSICAL MOTION IN FIELDS OF AHARONOV-BOHM AND AHARONOV-CASHER

Ya.I.Azimov^a, R.M.Ryndin^b
 Petersburg Nuclear Physics Institute
 Gatchina, St.Petersburg, 188350, Russia
e-mail: ^a azimov@pa1400.spb.edu,
^b ryndin@thd.pnpi.spb.ru

Abstract

Particle motion in the fields of Aharonov-Bohm and Aharonov-Casher is considered in framework of the classical theory to reveal conditions admitting duality of the two configurations. Important role of orientation of the magnetic dipole moment is demonstrated. Duality becomes totally destroyed by addition of electric dipole and/or higher multipole moments. Correspondence between quantum and classical considerations is also discussed.

PACS: 03.65.Bz

1. The Aharonov-Bohm effect (ABE) [1], the first example of purely topological effects closely related to the gauge character of electromagnetic field, continues to be of scientific and applied interest. Since its discovery the effect has been intensively studied in many details, but it was only rather recently that another topological effect in electromagnetic fields, analogous to ABE, was suggested, namely, the Aharonov-Casher effect (ACE) in purely electric field [2].

The question of existence of ACE initiated a theoretical discussion (see refs.[3, 4, 5]). It had not been concluded when the neutron beam measurements [6] confirmed ACE both qualitatively and with rather good quantitative

precision. Even more precise value for the Aharonov-Casher phase was obtained with molecular beams in the homogeneous electric field [7, 8]. And the theoretical discussion was closed by the paper of Hagen [9] who demonstrated exact duality of the Dirac wave functions in the fields of Aharonov-Bohm (AB) and Aharonov-Casher (AC).

The Hagen approach was later generalized for an arbitrary spin s , with the result [10]: the AB and AC configurations are mutually ‘dual only at the extreme spin projections, $\pm s$, normal to the field plane. Such result in the 2-dimensional problem seems to present a paradox: according to the group theory the total angular momentum should be inessential, its role should be played by the normal projection. For example, one could expect that the particle motion at spin $3/2$ and its normal projection $1/2$ should be equivalent to the motion at the same normal projection, but with spin $1/2$. However, the AB and AC configurations are dual for the latter case and not dual for the former one.

In the present note we show that purely classical consideration of particles in the fields of AB and AC reveals physical reasons for such special properties of the extreme spin projections. We can also study influence of higher multipole moments, above the charge and magnetic moment.

2. Begin with consideration of a purely classical particle in electric and magnetic fields. Comparison of papers [3, 4, 5] to each other and to earlier studies [11, 12] reveals absence of the generally accepted relativistic Lagrangian L_{int} for interaction of point-like multipoles with external fields. We construct it from two requirements:

- In the rest-frame of the particle (at $\mathbf{v} = 0$) L_{int} equals to the corresponding potential energy with sign reversed;
- L_{int} transforms as dS/dt with S being the invariant action, so it is an invariant multiplied by $(1 - v^2)^{1/2}$.

In this way we obtain, e.g., the following interaction Lagrangians for the point charge and the point-like electric and magnetic dipoles (such procedure gives, of course, the standard form to $L^{(ch)}$):

$$L^{(ch)}(\mathbf{x}, t) = -q\phi(\mathbf{x}, t) + q\mathbf{v} \cdot \mathbf{A}(\mathbf{x}, t) , \quad (1)$$

$$L^{(ed)}(\mathbf{x}, t) = \mathbf{d} \cdot \mathbf{E}(\mathbf{x}, t) + \mathbf{d} \cdot [\mathbf{v}\mathbf{B}(\mathbf{x}, t)] , \quad (2)$$

$$L^{(md)}(\mathbf{x}, t) = \mathbf{m} \cdot \mathbf{B}(\mathbf{x}, t) - \mathbf{m} \cdot [\mathbf{v}\mathbf{E}(\mathbf{x}, t)] . \quad (3)$$

There exists a tradition (see, e.g., refs.[11, 12]) to describe electric and magnetic dipoles by a combined antisymmetric tensor (similar to the field tensor $(\mathbf{E}, \mathbf{B})$). However, they play very different roles for elementary particles (e.g., $\mathbf{d} \neq 0$ leads to P and T violation). So we use two independent tensors, each one having a special form in the particle rest-frame: $(-\mathbf{d}_0, 0), (0, \mathbf{m}_0)$. In an arbitrary system these tensors are $(-\mathbf{d}(1-v^2)^{-1/2}, [\mathbf{d}\mathbf{v}](1-v^2)^{-1/2})$ and $([\mathbf{m}\mathbf{v}](1-v^2)^{-1/2}, \mathbf{m}(1-v^2)^{-1/2})$ if the particle moves with the velocity $\mathbf{v}$.

Correct relativistic transformation of L_{int} is provided by $q = \text{inv}$, while $\mathbf{d}$ and $\mathbf{m}$ should change. Transition from the rest-frame to a moving one transforms them just in the same way as the radius vector between two space points: longitudinal components undergo the Lorentz contraction, transversal ones do not change. Nevertheless, we do not vary them with $\mathbf{v}$: since dipole moments of a classical particle may change with time (in both value and direction) even in its rest-frame, we assume them to depend directly on time but not on velocity which is time-dependent itself.

The above interaction Lagrangians explicitly depend on velocity, so the particle canonical momentum contains both the familiar contribution, proportional to the vector potential, and other contributions, proportional to other multipoles. $L^{(ch)}$, $L^{(ed)}$ and $L^{(md)}$ produce additional terms

$$\delta \mathbf{p} = q_e \mathbf{A}(\mathbf{x}, t) - [\mathbf{d}\mathbf{B}(\mathbf{x}, t)] + [\mathbf{m}\mathbf{E}(\mathbf{x}, t)] . \quad (4)$$

The last two terms are sometimes considered as "hidden" momenta, but these extra terms are really induced by the explicit dependence of interaction on velocity.

Every multipole generates the corresponding forces of the particle interaction with fields. For $\mathbf{d}$ and $\mathbf{m}$ the forces may be presented as $\mathbf{f} = \mathbf{f}_{(0)} + \delta \mathbf{f}$, where

$$\mathbf{f}_{(0)}^{(ed)} = d_i \partial_i \mathbf{E} + d_i [\mathbf{v} \partial_i \mathbf{B}] + [\dot{\mathbf{d}}\mathbf{B}] , \quad (5)$$

$$\mathbf{f}_{(0)}^{(md)} = m_i \partial_i \mathbf{B} - m_i [\mathbf{v} \partial_i \mathbf{E}] - [\dot{\mathbf{m}}\mathbf{E}] . \quad (6)$$

Because of the Maxwell equations, the extra term $\delta \mathbf{f}^{(ed)}$ disappears while $\delta \mathbf{f}^{(md)}$ appears to be expressed through charge and current densities produced by the particle itself. So (if not vanishing) it corresponds to the particle self-interaction which influence should be included into the particle mass and/or other parameters, but should not arise in its equation of motion under external forces. Note that the forces $\mathbf{f}_{(0)}$ may be obtained by describing dipoles as limiting systems of two separated opposite charges, electric $\pm q_e$ or magnetic $\pm q_m$ (if they existed), each one subjected to the electric or magnetic Lorentz force

$$\mathbf{f}^{(e)} = q_e \mathbf{E} + q_e [\mathbf{v}\mathbf{B}] , \quad \mathbf{f}^{(m)} = q_m \mathbf{B} - q_m [\mathbf{v}\mathbf{E}] . \quad (7)$$

$\mathbf{f}^{(e)}$ has the familiar form and arises directly from $L^{(ch)}$; the both forces may be obtained by the Lorentz transformation of the force acting on a motionless charge, respectively electric or magnetic. Strange enough, none of papers [2, 3, 4, 5] on ACE gives the total classical force (6), though each of them correctly presents its separate terms.

There is one more point which has not been discussed at all in connection with ACE. A particle bearing dipole moments (or higher multipoles), even being point-like, should be considered as anisotropic. Its orientation produces new degrees of freedom. The particle can have non-vanishing angular momentum, it

can be subjected to torques. Expressions for the torques may be obtained from L_{int} by varying the particle orientation. For dipole moments the torques are

$$\mathbf{M}^{(ed)} = [\mathbf{d}\mathbf{E}] + [\mathbf{d}[\mathbf{v}\mathbf{B}]] , \quad (8)$$

$$\mathbf{M}^{(md)} = [\mathbf{m}\mathbf{B}] - [\mathbf{m}[\mathbf{v}\mathbf{E}]] . \quad (9)$$

We shall see in what follows that torques are very important for understanding properties of the particle motion.

In a similar way one can also find Lagrangians, forces and torques for particles with any higher multipoles. We do not give them here explicitly, but will have them in mind to consider the contribution of higher multipoles.

Let us briefly discuss how the motion influences interaction of multipoles with external fields. For L_{int} and $\mathbf{M}$ it leads only to a formal change of real fields by effective ones $\mathbf{E}' = \mathbf{E} + [\mathbf{v}\mathbf{B}]$, $\mathbf{B}' = \mathbf{B} - [\mathbf{v}\mathbf{E}]$. But the change is not the only physical effect, as can be seen from additional contributions to the canonical momentum and force.

3. Let us apply Lagrangians (1)-(3) to the classical motion of a particle in the Aharonov-Bohm (AB) field, i.e. in the field of the infinite thin solenoid along the z -axis. Field strengths vanish around the solenoid; the motion there looks free, without any forces and torques. However, $\mathbf{p}$ contains an extra term $\delta\mathbf{p}(\mathbf{x})$ with $\delta p_z = 0$.

The situation becomes less trivial if we consider the thin solenoid as the limit of a solenoid with a finite radius. For definiteness we suggest the following structure of the magnetic field strength: from the z -axis up to some radius the field is homogeneous and directed along the z -axis; in the transient region the field conserves its direction and diminishes its strength down to zero; the field strength (but not potential!) is absent everywhere outside. We also assume that initially the particle moves in the (x, y) plane orthogonal to the solenoid.

If the particle in its motion does not go through the strength region then no force and no torque arise; the classical motion stays free. In the strength region, as is well known, the field produces the force $\mathbf{f}^{(e)}$. It acts on the particle charge and tends to curve its trajectory without driving it out of the (x, y) -plane. Neither the particle orientation changes.

Let us consider the role of magnetic moment. If the dipole is normal to the motion plane (i.e. along the z -axis) then, according to (6) and (9), it is not subjected to any torque or additional force. The particle orientation conserves.

The situation is quite different if $\mathbf{m}$ has non-vanishing projection onto the motion plane. Because of the torque $\mathbf{M}^{(md)}$ the dipole $\mathbf{m}$ precesses around the field direction. Precession does not produce any additional force in the homogeneous region. However, in the transient region it does generate the force $\mathbf{f}^{(md)}$ which has a non-vanishing z -component and draws the motion out of the (x, y) -plane. The sign of $f_z^{(md)}$ may change in the course of precession, so the motion oscillates around its initial plane. Thus, the particle leaves the field region with, generally, non-zero v_z . Its sign and value depend on both the field

structure in the transient region and the initial orientation of $\mathbf{m}$. Projection of $\mathbf{m}$ onto the $\mathbf{B}$ -direction does not change.

The motion becomes much more complicated if the particle, in addition to the charge and magnetic moment, has also an electric dipole moment (EDM) or higher multipoles. In such a case the precession axis itself changes its direction, and the particle orientation evolves in a rather complicated way. There are additional forces and torques acting in both transient and homogeneous regions, at any initial particle orientation. As a result, after going through the field the particle not only gets out of plane, but also changes its $\mathbf{m}$ -projection onto $\mathbf{B}$.

So, we see that a classical particle in the field of a long solenoid moves differently in the presence, in addition to the charge, of some other multipoles (even magnetic moment). Resulting motion is very sensitive to the field structure, and, hence, the problem of the infinitely long and thin solenoid may essentially depend on the limiting procedure. Only for the particle with charge and magnetic moment, exactly along the field direction, the motion has "canonical" properties.

4. Now we consider a classical particle in the Aharonov-Casher (AC) field, i.e., in the field of a uniformly charged straight thin thread along the z -axis. Its electric strength is perpendicular to the thread and depends only on two coordinates x and y . We assume the particle to be neutral, but having a magnetic moment and, may be, some other multipoles. Initially the particle moves in the (x, y) -plane.

Let us begin with $\mathbf{m}$ oriented along the z -axis. Then eq.(9) shows that torques are absent, and the particle orientation does not change. Furthermore, $\dot{\mathbf{m}} = 0$, $m_x = m_y = 0$, and, according to eq.(6), forces are also absent. The particle motion looks to be free, but its canonical momentum, according to eq.(4), differs from the free value $m\mathbf{v}(1 - v^2)^{-1/2}$. The extra term $\delta\mathbf{p}$ has the same structure as for the charged particle in the AB field outside the solenoid (in particular, $\delta p_z = 0$). Therefore, all the classical description of this case is similar to the case of a charged particle in the AB field.

If $\mathbf{m}$ has non-vanishing projection onto the (x, y) -plane, then $\delta p_z \neq 0$. This means that $\delta\mathbf{p}$ is definitely different here from the extra term in the AB case. Such motion generates the torque which compels $\mathbf{m}$ to precess around the axis $[\mathbf{v}\mathbf{E}]$, initially parallel to the z -direction. Therefore, non-vanishing $\dot{\mathbf{m}}$ arises, initially in the (x, y) -plane. Note that $\dot{\mathbf{m}}$ and $\mathbf{m}$ are orthogonal to each other, so the absolute value $|\mathbf{m}|$ does not change. Now, non-zero $\dot{\mathbf{m}}$ and projections of $\mathbf{m}$ onto the (x, y) -plane generate, according to (6), a force (with non-zero z -component) which changes the $\mathbf{v}$ -direction and even draws the motion out of the plane.

If one compares such classical motion in the AC configuration to the classical motion of a charged particle in the AB field with the same initial orientation of $\mathbf{m}$, then an essential difference can be seen. The AB motion is also subjected to forces and torques violating its plane character, but only when propagating through the confined region of the solenoid itself. In the AC field the forces and

torques act over the whole space. Thus, the AC motion with oblique orientation of $\mathbf{m}$ can never be free and never is similar to the AB motion. The magnetic moment $\mathbf{m}$ conserves its absolute value, but changes its orientation in a rather complicated way: it precesses around the axis $[\mathbf{v}\mathbf{E}]$ which itself changes its direction.

Nearly of the same character is the motion when the particle carries also EDM or higher multipoles. But time evolution of $\mathbf{m}$ and $\mathbf{v}$ becomes more complicated. One of the reasons is that addition of any other multipoles to the magnetic moment induces forces and torques which affect the motion in the AC field even for the initial $\mathbf{m}$ -orientation along the z -axis. Therefore, possibility of duality between the classical AB and AC motions becomes totally destroyed.

5. In conclusion, we briefly discuss correspondence between classical and quantum descriptions of multipole interactions for a moving particle. We have shown in the preceding Sections that similarity between the classical AB and AC motions is possible only if the magnetic moment was initially oriented exactly along the z -axis. Such a conclusion is just the same as the result of the quantum consideration [10] and reveals its physical reason.

Furthermore, the classical study has shown that addition of EDM or higher multipoles totally prevents possible AB–AC duality. Leaving apart detailed quantum analysis of such a case we consider here only briefly how the multipoles influence the particle wave function.

In the AB field the particle carrying only charge and magnetic moment conserves its spin z -projection [9, 10]. This property for a Dirac particle in the AB field is expressed by commutation of $\gamma_3\gamma_5$ with the Dirac operator, if there is only magnetic field directed along the z -axis and the wave function depends only on x and y . Presence of EDM induces interaction proportional to $\sigma_{\mu\nu}\gamma_5 F^{\mu\nu}$ (P and T violation does not matter here). If the field consists only of B_z , we have the operator contribution $\sigma_{12}\gamma_5$ which does not commute with $\gamma_3\gamma_5$, and the spin orientation cannot conserve. EDM has similar effect in the AC configuration, since it induces the terms $\sigma_{0k}\gamma_5$ with $k = 1, 2$ in the Dirac operator. However, the quantitative influence of EDM on the AB and AC wave functions is different, both because of different properties of fields $\mathbf{B}$ and $\mathbf{E}$ and due to their different space distributions. So, the AB–AC duality is excluded by the presence of EDM.

Higher multipoles require a higher spin and complicate relativistic consideration (see, e.g., [10]). But for the qualitative understanding we may stick to the quadrupole moment in nonrelativistic approximation. Operator of the quadrupole moment, both electric and magnetic, is proportional to $S_j S_k + S_k S_j - \frac{2}{3}\mathbf{S}^2\delta_{jk}$, where $\mathbf{S}$ is the spin operator. It is easy to check that in both configurations, AB and AC, the interaction contains terms not commuting with S_3 and violating the spin z -component conservation. The AB–AC duality is again impossible. So, the quantum description of the role of various multipoles also agrees with classical one, at least qualitatively.

One of interesting classical results is the inevitable non-plane character of

motion if the magnetic moment was initially deflected from the z -axis (or in the presence of EDM or higher multipoles). Meanwhile, the quantum wave function may still depend on x and y only which is usually thought to indicate plane motion. However, really it is not so. At the space infinity, where $\mathbf{E}$ vanishes, the canonical momentum coincides with $m\mathbf{v}(1 - v^2)^{-1/2}$. Let us assume that $v_z = 0$ there and so $\mathbf{p}$ lies in the (x, y) -plane. In the AC motion p_z conserves and $\mathbf{p}$ remains in the plane. That is why the wave function depends only on two variables. But, if $\mathbf{m}$ was deflected from the z -axis, $\mathbf{v}$ deviates from $\mathbf{p}$ (see eq.(4)) and acquires non-zero z -component. Only at $|m_z| = |\mathbf{m}|$ the problem degenerates into the really plane one. Detailed comparison of quantum and classical considerations here may require to define more precisely what is the velocity (and, m.b., coordinate as well) for a quantum relativistic particle in the external field.

As a result of classical consideration and its comparison to quantum one we can note an interesting difference between ABE and ACE. In the field of a thin solenoid the wave function of the charged particle, while going around the solenoid, produces the known geometrical phase independently of the form of the solenoid and orientation of the particle magnetic moment (the solenoid is straight only for simplicity; particle orientation is not important since the field strength vanishes around the solenoid). On the contrary, orientation is very essential in the field of a thin charged thread: even for the straight thread the geometrical phase arises only for two extreme orientations of the magnetic moment. But if the thread is curved, the magnetic moment projection onto the thread direction cannot be fixed, and going around the thread is not related to the geometrical phase. Therefore, ACE disappears if the charged thread is bent, while the ABE is just a direct consequence of the presence of the solenoid, independently of its form. In that sense, ACE is not a real topological effect.

This work has been supported by the grant RFBR 96-02-18630. The authors are grateful to A.Vainshtein and V.Zelevinsky for stimulating discussions.

References

- [1] Y.Aharonov, D.Bohm, Phys.Rev. **115**, 485 (1959).
- [2] Y.Aharonov, A.Casher, Phys.Rev.Lett. **53**, 319 (1984).
- [3] T.H.Boyer, Phys.Rev. **A36**, 5083 (1987).
- [4] Y.Aharonov, P.Pearle, L.Vaidman, Phys.Rev. **A37**, 4052 (1988).
- [5] A.S.Goldhaber, Phys.Rev.Lett. **62**, 482 (1989).
- [6] A.Cimmino et al., Phys.Rev.Lett. **63**, 380 (1989).
- [7] K.Sangster, E.A.Hinds, S.M.Barnett, E.Riis, Phys.Rev.Lett. **71**, 3641 (1993).

- [8] K.Sangster, E.A.Hinds, S.M.Barnett et al., Phys.Rev. **A51**, 1776 (1995).
- [9] C.R.Hagen, Phys.Rev.Lett. **64**, 503 (1990).
- [10] Ya.I.Azimov, R.M.Ryndin, Pis'ma ZhETF **61**, 444 (1995) [JETP Letters **61**, 453 (1995)].
- [11] W.Pauli. Theory of relativity. London, 1958.
- [12] Ya.I.Frenkel. Electrodynamics. ONTI publishers, Moscow, 1934 (in Russian).